

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/SE05/000448

International filing date: 29 March 2005 (29.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: SE
Number: 0400850-4
Filing date: 30 March 2004 (30.03.2004)

Date of receipt at the International Bureau: 29 April 2005 (29.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

PRV

PATENT- OCH REGISTRERINGSVERKET

Patentavdelningen

**Intyg
Certificate**

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande AstraZeneca AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 0400850-4
Patent application number

(86) Ingivningsdatum 2004-03-30
Date of filing

Stockholm, 2005-04-06

*För Patent- och registreringsverket
For the Patent- and Registration Office*

Gunilla Larsson
Gunilla Larsson

Avgift
Fee

The present invention relates to novel triazolone derivatives, processes for their preparation, pharmaceutical compositions containing them and their use in therapy.

5 Metalloproteinases are a superfamily of proteinases (enzymes) whose numbers in recent years have increased dramatically. Based on structural and functional considerations these enzymes have been classified into families and subfamilies as described in N.M. Hooper (1994) FEBS Letters 354:1-6. Examples of metalloproteinases include the matrix metalloproteinases (MMPs) such as the collagenases (MMP1, MMP8, MMP13), the gelatinases (MMP2, MMP9), the stromelysins (MMP3, MMP10, MMP11), matrilysin 10 (MMP7), metalloelastase (MMP12), enamelysin (MMP19), the MT-MMPs (MMP14, MMP15, MMP16, MMP17); the reprolysin or adamalysin or MDC family which includes the secretases and sheddases such as TNF converting enzymes (ADAM10 and TACE); the astacin family which include enzymes such as procollagen processing proteinase (PCP); and other metalloproteinases such as aggrecanase, the endothelin converting enzyme 15 family and the angiotensin converting enzyme family.

Metalloproteinases are believed to be important in a plethora of physiological disease processes that involve tissue remodelling such as embryonic development, bone formation and uterine remodelling during menstruation. This is based on the ability of the 20 metalloproteinases to cleave a broad range of matrix substrates such as collagen, proteoglycan and fibronectin. Metalloproteinases are also believed to be important in the processing, or secretion, of biological important cell mediators, such as tumour necrosis factor (TNF); and the post translational proteolysis processing, or shedding, of biologically important membrane proteins, such as the low affinity IgE receptor CD23 (for a more 25 complete list see N. M. Hooper *et al.*, (1997) Biochem J. 321:265-279).

Metalloproteinases have been associated with many diseases or conditions. Inhibition of the activity of one or more metalloproteinases may well be of benefit in these diseases or conditions, for example: various inflammatory and allergic diseases such as, inflammation

of the joint (especially rheumatoid arthritis, osteoarthritis and gout), inflammation of the gastro-intestinal tract (especially inflammatory bowel disease, ulcerative colitis and gastritis), inflammation of the skin (especially psoriasis, eczema, dermatitis); in tumour metastasis or invasion; in disease associated with uncontrolled degradation of the
5 extracellular matrix such as osteoarthritis; in bone resorptive disease (such as osteoporosis and Paget's disease); in diseases associated with aberrant angiogenesis; the enhanced collagen remodelling associated with diabetes, periodontal disease (such as gingivitis), corneal ulceration, ulceration of the skin, post-operative conditions (such as colonic anastomosis) and dermal wound healing; demyelinating diseases of the central and
10 peripheral nervous systems (such as multiple sclerosis); Alzheimer's disease; extracellular matrix remodelling observed in cardiovascular diseases such as restenosis and atherosclerosis; asthma; rhinitis; and chronic obstructive pulmonary diseases (COPD).

MMP12, also known as macrophage elastase or metalloelastase, was initially cloned in the
15 mouse by Shapiro *et al* [1992, Journal of Biological Chemistry 267: 4664] and in man by the same group in 1995. MMP12 is preferentially expressed in activated macrophages, and has been shown to be secreted from alveolar macrophages from smokers [Shapiro *et al*, 1993, Journal of Biological Chemistry, 268: 23824] as well as in foam cells in atherosclerotic lesions [Matsumoto *et al*, 1998, Am J Pathol 153: 109]. A mouse model of
20 COPD is based on challenge of mice with cigarette smoke for six months, two cigarettes a day six days a week. Wild-type mice developed pulmonary emphysema after this treatment. When MMP12 knock-out mice were tested in this model they developed no significant emphysema, strongly indicating that MMP12 is a key enzyme in the COPD pathogenesis. The role of MMPs such as MMP12 in COPD (emphysema and bronchitis) is
25 discussed in Anderson and Shinagawa, 1999, Current Opinion in Anti-inflammatory and Immunomodulatory Investigational Drugs 1(1): 29-38. It was recently discovered that smoking increases macrophage infiltration and macrophage-derived MMP-12 expression in human carotid artery plaques Kangavari [Matetzky S, Fishbein MC *et al.*, Circulation 102:(18), 36-39 Suppl. S, Oct 31, 2000].

MMP9 (Gelatinase B; 92kDa TypeIV Collagenase; 92kDa Gelatinase) is a secreted protein which was first purified, then cloned and sequenced, in 1989 [S.M. Wilhelm *et al* (1989) J. Biol Chem. 264 (29): 17213-17221; published erratum in J. Biol Chem. (1990) 265 (36): 22570]. A recent review of MMP9 provides an excellent source for detailed information and references on this protease: T.H. Vu & Z. Werb (1998) (In : Matrix Metalloproteinases. 1998. Edited by W.C. Parks & R.P. Mecham. pp115 - 148. Academic Press. ISBN 0-12-545090-7). The following points are drawn from that review by T.H. Vu & Z. Werb (1998).

The expression of MMP9 is restricted normally to a few cell types, including trophoblasts, osteoclasts, neutrophils and macrophages. However, its expression can be induced in these same cells and in other cell types by several mediators, including exposure of the cells to growth factors or cytokines. These are the same mediators often implicated in initiating an inflammatory response. As with other secreted MMPs, MMP9 is released as an inactive Pro-enzyme which is subsequently cleaved to form the enzymatically active enzyme. The proteases required for this activation *in vivo* are not known. The balance of active MMP9 versus inactive enzyme is further regulated *in vivo* by interaction with TIMP-1 (Tissue Inhibitor of Metalloproteinases -1), a naturally-occurring protein. TIMP-1 binds to the C-terminal region of MMP9, leading to inhibition of the catalytic domain of MMP9. The balance of induced expression of ProMMP9, cleavage of Pro- to active MMP9 and the presence of TIMP-1 combine to determine the amount of catalytically active MMP9 which is present at a local site. Proteolytically active MMP9 attacks substrates which include gelatin, elastin, and native Type IV and Type V collagens; it has no activity against native Type I collagen, proteoglycans or laminins.

There has been a growing body of data implicating roles for MMP9 in various physiological and pathological processes. Physiological roles include the invasion of embryonic trophoblasts through the uterine epithelium in the early stages of embryonic

implantation; some role in the growth and development of bones; and migration of inflammatory cells from the vasculature into tissues.


5 MMP9 release, measured using enzyme immunoassay, was significantly enhanced in fluids and in AM supernatants from untreated asthmatics compared with those from other populations [Am. J. Resp. Cell & Mol. Biol., Nov 1997, 17 (5):583-591]. Also, increased MMP9 expression has been observed in certain other pathological conditions, thereby implicating MMP9 in disease processes such as COPD, arthritis, tumour metastasis, Alzheimer's, Multiple Sclerosis, and plaque rupture in atherosclerosis leading to acute
10 coronary conditions such as Myocardial Infarction.

A number of metalloproteinase inhibitors are known (see for example the reviews of MMP inhibitors by Beckett R.P. and Whittaker M., 1998, Exp. Opin. Ther. Patents, 8(3):259-282, and by Whittaker M. *et al*, 1999, Chemical Reviews 99(9):2735-2776).

15 We have now discovered a new class of compounds, namely triazolone derivatives, that are inhibitors of metalloproteinases and are of particular interest in inhibiting MMPs such as MMP12 and MMP9. The compounds of the present invention have beneficial potency, selectivity and/or pharmacokinetic properties. Certain compounds of the invention may
20 also be useful as inhibitors of TACE and/or aggrecanase.

In accordance with the present invention, there is therefore provided a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof

5



(I)

wherein

5 R^1 and R^2 independently represent H or C1 to 6 alkyl; said alkyl being optionally further substituted by an aryl ring or an aromatic heterocyclic ring containing 1 to 3 heteroatoms independently selected from O, S and N; said aromatic ring being optionally further substituted by halogen, CF_3 , C1 to 4 alkyl or C1 to 4 alkoxy;

10 Each R^3 and each R^4 independently represents H or C1 to 6 alkyl; said alkyl being optionally further substituted by OH, C1 to 4 alkoxy, C1 to 4 alkylthio, amino, N-alkylamino or N,N-dialkylamino;

15 or R^3 and R^4 are bonded together so as to form a 3 to 7 membered ring; said ring optionally incorporating one heteroatom selected from O, $S(O)_q$ and N;

m represents an integer 1, 2 or 3;

20 X represents a group $S(O)$, $S(O)_2$ or $C(=O)$;

R^5 represents H or C1 to 6 alkyl; said alkyl being optionally further substituted by halogen, OH or C1 to 6 alkoxy;

Y represents a direct bond;

5

or Y and R^5 are bonded together such that the group $-NR^5Y-$ together represents a 4 to 7 membered saturated or partially unsaturated azacyclic ring; said azacyclic ring optionally incorporating one further heteroatom selected from O, $S(O)_n$ and N; said azacyclic ring being optionally benzo fused; said azacyclic ring being optionally substituted by C1 to 6 alkyl, C1 to 6 alkoxy or OH;

10

L represents a direct bond;

or L represents O, $S(O)_p$, $C(O)$, NR^6 , $C(O)NR^6$, $NR^6C(O)$, C2 to 6 alkynyl, C2 to 6 alkenyl, C1 to 6 alkyl, C1 to 6 heteroalkyl or C3 to 6 heteroalkynyl; said alkyl, alkenyl or alkynyl group being optionally further substituted by halogen, OH or C1 to 6 alkoxy;

15

n, p and q independently represent an integer 0, 1 or 2;

G^1 represents a monocyclic, bicyclic, tricyclic or tetracyclic group comprising one, two, three or four ring structures each of up to 7 ring atoms; each ring structure being independently selected from cycloalkyl; cycloalkenyl; heterocycloalkyl; unsaturated heterocycloalkyl; aryl; or an aromatic heterocyclic ring containing 1 to 3 heteroatoms independently selected from O, S and N; with each ring structure being independently optionally substituted by one or more substituents independently selected from halogen, hydroxy, CHO, C1 to 6 alkyl, C1 to 6 alkoxy, halo-C1 to 6 alkoxy, amino, N-alkylamino, N,N-dialkylamino, alkylsulfonamino, C2 to 6 alkanoylamino, cyano, nitro, thiol, alkylthio, alkylsulfonyl, alkylaminosulfonyl, C2 to 6 alkanoyl, aminocarbonyl, N-alkylamino-carbonyl, N,N-amino-carbonyl;

25

wherein any alkyl radical within any substituent may itself be optionally substituted with one or more groups selected from halogen, hydroxy, C1 to 6 alkoxy, halo-C1 to 6 alkoxy, amino, N-alkylamino, N,N-dialkylamino, N-alkylsulfonamino, N-C2 to 6 alkanoylamino, cyano, nitro, thiol, alkylthio, alkylsulfonyl, N-alkylaminosulfonyl, CHO, C2 to 6 alkanoyl, aminocarbonyl, N-alkylaminocarbonyl, N,N-dialkylaminocarbonyl and carbamate;

and wherein any alkyl radical is a C1 to 6 alkyl radical;

and when G^1 is a bicyclic, tricyclic or tetracyclic group, each ring structure is independently joined to the next ring structure by a direct bond, by -O-, by C1-6 alkyl, by C1-6 haloalkyl, by C1-6 heteroalkyl, by C2-6 alkenyl, by C2-6 alkynyl, by sulfone, by CO, by NR^7CO , by $CONR^7$, by NR^7 , by S, or by C(OH), or each ring structure is fused to the next ring structure;


R^6 and R^7 independently represent H or C1 to 6 alkyl;

and when the group $-NR^5Y-$ represents an azacyclic ring and L represents a direct bond, the group G^1 may also be spiro fused to the azacyclic ring;

and pharmaceutically acceptable salts thereof.

The compounds of formula (I) may exist in enantiomeric forms. It is to be understood that all enantiomers, diastereomers, racemates and mixtures thereof are included within the scope of the invention.

Compounds of formula (I) may also exist in various tautomeric forms. Thus, for example, the triazolone ring of compounds in which R^1 and R^2 each represent H can exist in the following tautomeric forms:



All possible tautomeric forms and mixtures thereof are included within the scope of the invention.

- 10 In one embodiment, X represents $S(O)_2$. In another embodiment, X represents $C(=O)$.

In one embodiment, R^1 represents H. In one embodiment, R^2 represents H. In another embodiment, R^1 and R^2 each represent H.

- 15 In one embodiment, R^3 and R^4 independently represent H or C1 to 6 alkyl. In another embodiment, R^3 and R^4 each represent H.

In one embodiment, m represents the integer 1. In another embodiment, m represents the integer 2.

20

In one embodiment, R^5 represents H or C1 to 6 alkyl. In another embodiment, R^5 represents H.

In one embodiment, Y represents a direct bond.

25

In another embodiment, Y and R^5 are bonded together such that the group $-NR^5Y-$ together represents a 4 to 7 membered saturated or partially unsaturated azacyclic ring;

said azacyclic ring optionally containing one further heteroatom selected from O, S(O)_n and N; said azacyclic ring being optionally benzo fused.

In another embodiment, Y and R⁵ are bonded together such that the group -NR⁵Y-
5 together represents a 4 to 7 membered saturated or partially unsaturated azacyclic ring;
said azacyclic ring optionally containing one further heteroatom selected from O, S(O)_n
and N.

In another embodiment, Y and R⁵ are bonded together such that the group -NR⁵Y-
10 together represents piperidinyl, 3,4-dehydropiperidinyl or piperazinyl.

In one embodiment, L represents a direct bond. In another embodiment, L represents O, C2
to 6 alkynyl, C1 to 6 alkyl, C1 to 6 heteroalkyl or C3 to 6 heteroalkynyl.

15 In one embodiment, G¹ represents an optionally substituted monocyclic or bicyclic ring
structure. In another embodiment, G¹ represents an optionally substituted monocyclic ring
structure. In another embodiment, G¹ represents an optionally substituted phenyl or
heteroaryl ring. In another embodiment, G¹ represents an optionally substituted bicyclic
ring structure. In another embodiment, G¹ represents an optionally substituted bicyclic ring
20 structure in which each ring is independently phenyl or heteroaryl. In another
embodiment, G¹ represents an optionally substituted bicyclic ring structure in which the
two rings are either bonded directly to one another or are separated by an O atom. In
another embodiment, G¹ represents an optionally substituted bicyclic ring structure in
which each ring is independently phenyl or heteroaryl and the two rings are either bonded
25 directly to one another or are separated by an O atom.

Unless otherwise indicated, the term "C1 to 6 alkyl" referred to herein denotes a straight or
branched chain alkyl group having from 1 to 6 carbon atoms. Examples of such groups

include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl and t-butyl. The term "C1 to 4 alkyl" is to be interpreted analogously.

The two alkyl moieties in a dialkylamino group may be the same or different.

Unless otherwise indicated, the term "C2 to 6 alkenyl" referred to herein denotes a straight or branched chain alkyl group having from 2 to 6 carbon atoms incorporating at least one carbon-carbon double bond. Examples of such groups include ethenyl, propenyl and butenyl.

Unless otherwise indicated, the term "C2 to 6 alkynyl" referred to herein denotes a straight or branched chain alkyl group having from 2 to 6 carbon atoms incorporating at least one carbon-carbon triple bond. Examples of such groups include ethynyl, propynyl, and butynyl.

Unless otherwise indicated, the term "C1 to 6 alkoxy" referred to herein denotes a straight or branched chain alkyl group having from 1 to 6 carbon atoms bonded to a molecule via an oxygen atom. Examples of such groups include methoxy, ethoxy, n-propoxy, i-propoxy and t-butoxy. The term "C1 to 6 alkylthio" is to be interpreted analogously but with bonding being via a sulphur atom. The terms "C1 to 4 alkoxy" and "C1 to 4 alkylthio" are to be interpreted analogously.

Unless otherwise indicated, the term "halogen" referred to herein denotes fluoro, chloro, bromo and iodo.

Unless otherwise indicated, the term "C1 to 6 heteroalkyl" referred to herein denotes a straight or branched chain alkyl group having from 1 to 6 carbon atoms and incorporating one or more heteroatoms selected independently from O, S(O)_n and N. Examples of such groups include -O-(CH₂)₃-, -CH₂CH₂OCH₂-, -CH₂CH₂SCH₂CH₂-,

$-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{OCH}_2-$. The term "C3 to 6 heteroalkynyl" is to be interpreted analogously and would include such groups as $-\text{C}\equiv\text{C}-\text{CH}_2-\text{O}-$.

Examples of a "C1 to 6 haloalkyl or halo-C1 to 6 alkoxy" include CH_2F , CHF_2 , CF_3 ,
5 CF_3CF_2 , CF_3CH_2 , CH_2FCH_2 , CH_3CF_2 , $\text{CF}_3\text{CH}_2\text{CH}_2$, OCF_3 and OCH_2CF_3 .

Unless otherwise indicated, the term "C2 to 6 alkanoyl" referred to herein denotes a straight or branched chain alkyl group having from 1 to 5 carbon atoms bonded to a molecule via a carbonyl ($\text{C}=\text{O}$) group. Examples of such groups include acetyl, propionyl
10 and pivaloyl.

Examples of a 4 to 7 membered saturated or partially unsaturated azacyclic ring; optionally incorporating one further heteroatom selected from O, $\text{S}(\text{O})_n$ or N; and optionally being benzo fused; include pyrrolidine, piperidine, 3,4-dehydropiperidine, tetrahydroquinoline,
15 tetrahydrosioquinoline, piperazine, morpholine and perhydroazepine.

Examples of an aromatic heterocyclic ring of up to 7 ring atoms containing 1 to 3 heteroatoms independently selected from O, S and N include furan, thiophene, pyrrole, pyridine, thiazole, imidazole, oxazole, isoxazole, pyrazole, triazole, oxadiazole,
20 thiadiazole, pyrazine, pyridazine and pyrimidine.



Examples of a cycloalkyl or cycloalkenyl ring containing up to 7 ring atoms include cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl and cyclohexenyl.

25 Examples of a heterocycloalkyl or unsaturated heterocycloalkyl ring containing up to 7 ring atoms include pyrrolidine, tetrahydrofuran, dioxane, dioxolane, thiane, piperidine, 3,4-dehydropiperidine, piperazine, morpholine, thiomorpholine and perhydroazepine.

Examples of an aryl group include phenyl and naphthyl.


Examples of compounds wherein the group $-NR^5Y-$ represents an azacyclic ring and L represents a direct bond and the group G^1 is spiro fused to the azacyclic ring include structures such as:

5




Specific examples of the molecular fragment

10



include



and corresponding structures in which the various rings are optionally substituted.

- 5 Specific examples of fused bicyclic ring systems include quinolinyl, isoquinolinyl, indolyl, tetrahydroisoquinolinyl, benzofuranyl, benzothienyl, quinazolinyl, phthalazinyl, dihydrobenzofuranyl, naphthyl and dihydroindolyl. Preferred bicyclic ring systems

include quinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, naphthyl, benzofuranyl and benzothienyl.

It will be appreciated that the particular substituents and number of substituents in the compounds of the invention are selected so as to avoid sterically undesirable combinations.

Examples of compounds of the invention include:

- 5-[(4-[(5-chloropyridin-2-yl)oxy]piperidin-1-yl)sulfonyl)methyl]-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-[2-[(4-[(5-chloropyridin-2-yl)oxy]piperidin-1-yl)sulfonyl]ethyl]-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-[3-[(4-[(5-chloropyridin-2-yl)oxy]piperidin-1-yl)sulfonyl]propyl]-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-([4-(4-chlorophenyl)piperazin-1-yl]sulfonyl)methyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-(2-[4-(4-chlorophenyl)piperidin-1-yl]sulfonyl)ethyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-(2-[4-(4-chlorophenyl)piperazin-1-yl]sulfonyl)ethyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-(3-[4-(4-chlorophenyl)piperidin-1-yl]sulfonyl)propyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-(3-[4-(4-chlorophenyl)piperazin-1-yl]sulfonyl)propyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-([4-[(2-methoxypyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2H)-yl]sulfonyl)methyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;
- and pharmaceutically acceptable salts and solvates thereof.

Each exemplified compound represents a particular and independent aspect of the invention.

The compounds of formula (I) may exist in enantiomeric forms. Therefore, all enantiomers, diastereomers, racemates and mixtures thereof are included within the scope of the invention. The various optical isomers may be isolated by separation of a racemic mixture of the compounds using conventional techniques, for example, fractional crystallisation, or HPLC.

- 5 Alternatively the optical isomers may be obtained by asymmetric synthesis, or by synthesis from optically active starting materials.

Where optically isomers exist in the compounds of the invention, we disclose all individual optically active forms and combinations of these as individual specific embodiments of the invention, as well as their corresponding racemates.

Where tautomers exist in the compounds of the invention, we disclose all individual tautomeric forms and combinations of these as individual specific embodiments of the invention.

15

The present invention includes compounds of formula (I) in the form of salts. Suitable salts include those formed with organic or inorganic acids or organic or inorganic bases. Such salts will normally be pharmaceutically acceptable salts although non-pharmaceutically acceptable salts may be of utility in the preparation and purification of particular compounds. Such salts include acid addition salts such as hydrochloride, hydrobromide, citrate, tosylate and maleate salts and salts formed with phosphoric acid or sulphuric acid. In another aspect suitable salts are base salts such as an alkali metal salt, for example, sodium or potassium, an alkaline earth metal salt, for example, calcium or magnesium, or an organic amine salt, for example, triethylamine. Examples of solvates include hydrates.

25

Salts of compounds of formula (I) may be formed by reacting the free base or another salt thereof with one or more equivalents of an appropriate acid or base.

The compounds of formula (I) are useful because they possess pharmacological activity in animals and are thus potentially useful as pharmaceuticals. In particular, the compounds of the invention are metalloproteinase inhibitors and may thus be used in the treatment of diseases or conditions mediated by MMP12 and/or MMP9 such as asthma, rhinitis, chronic
5 obstructive pulmonary diseases (COPD), arthritis (such as rheumatoid arthritis and osteoarthritis), atherosclerosis and restenosis, cancer, invasion and metastasis, diseases involving tissue destruction, loosening of hip joint replacements, periodontal disease, fibrotic disease, infarction and heart disease, liver and renal fibrosis, endometriosis, diseases related to the weakening of the extracellular matrix, heart failure, aortic
10 aneurysms, CNS related diseases such as Alzheimer's disease and Multiple Sclerosis (MS), and hematological disorders.

Accordingly, the present invention provides a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined for use in
15 therapy.

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

20

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in the treatment of diseases or conditions in which inhibition of MMP12 and/or MMP9 is beneficial.

25

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in the treatment of inflammatory disease.

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in the treatment of an obstructive airways disease such as asthma or COPD.

5

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

- 10 Prophylaxis is expected to be particularly relevant to the treatment of persons who have suffered a previous episode of, or are otherwise considered to be at increased risk of, the disease or condition in question. Persons at risk of developing a particular disease or condition generally include those having a family history of the disease or condition, or those who have been identified by genetic testing or screening to be particularly
- 15 susceptible to developing the disease or condition.

The invention further provides a method of treating a disease or condition in which inhibition of MMP12 and/or MMP9 is beneficial which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or a

20 pharmaceutically acceptable salt or solvate thereof as hereinbefore defined.

The invention also provides a method of treating an obstructive airways disease, for example, asthma or COPD, which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or

25 solvate thereof as hereinbefore defined.

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder to be treated. The daily dosage of the compound of formula (I)/salt/solvate (active

ingredient) may be in the range from 0.001 mg/kg to 75 mg/kg, in particular from 0.5 mg/kg to 30 mg/kg. This daily dose may be given in divided doses as necessary. Typically unit dosage forms will contain about 1 mg to 500 mg of a compound of this invention.

5

The compounds of formula (I) and pharmaceutically acceptable salts and solvates thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (I) compound/salt/solvate (active ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.10 to 70 %w, of active ingredient, and, from 1 to 99.95 %w, more preferably from 30 to 99.90 %w, of a pharmaceutically acceptable adjuvant, diluent or carrier, all percentages by weight being based on total composition. Conventional procedures for the selection and preparation of suitable pharmaceutical formulations are described in, for example, "Pharmaceuticals - The Science of Dosage Form Designs", M. E. Aulton, Churchill Livingstone, 1988.

15

Thus, the present invention also provides a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as hereinbefore defined in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

20

The invention further provides a process for the preparation of a pharmaceutical composition of the invention which comprises mixing a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as hereinbefore defined with a pharmaceutically acceptable adjuvant, diluent or carrier.


25

The pharmaceutical compositions of this invention may be administered in standard manner for the disease or condition that it is desired to treat, for example by oral, topical,


parenteral, buccal, nasal, vaginal or rectal administration or by inhalation. For these purposes the compounds of this invention may be formulated by means known in the art into the form of, for example, tablets, capsules, aqueous or oily solutions, suspensions, emulsions, creams, ointments, gels, nasal sprays, suppositories, finely divided powders or aerosols for inhalation, and for parenteral use (including intravenous, intramuscular or infusion) sterile aqueous or oily solutions or suspensions or sterile emulsions.

In addition to the compounds of the present invention the pharmaceutical composition of this invention may also contain, or be co-administered (simultaneously or sequentially) with, one or more pharmacological agents of value in treating one or more diseases or conditions referred to hereinabove such as "Symbicort" (trade mark) product.

The present invention further provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as defined above which comprises:
reaction of a compound of formula (II)



wherein R^1 , R^2 , R^3 , R^4 , X and m are as defined in formula (I) and L^1 represents a leaving group, with a compound of formula (III)



(III)

wherein G^1 , L , Y and R^5 are as defined in formula (I)

and optionally thereafter forming a pharmaceutically acceptable salt or solvate.

5

In the above process, suitable leaving groups L^1 include halo, particularly chloro. The reaction is preferably performed in a suitable solvent optionally in the presence of an added base for a suitable period of time, typically 1 to 24 h, at ambient to reflux temperature. Preferably, solvents such as pyridine, dimethylformamide, tetrahydrofuran, acetonitrile or dichloromethane are used. When used the added base may be an organic base such as triethylamine, diisopropylethylamine, N-methylmorpholine or pyridine, or an inorganic base such as an alkali metal carbonate. The reaction is typically conducted at ambient temperature for 2 to 16 h, or until completion of the reaction has been achieved, as determined by chromatographic or spectroscopic methods. Reactions of sulfonyl halides and acyl halides with various primary and secondary amines are well known in the literature, and the variations of the conditions will be evident for those skilled in the art.


10

15

Compounds of formula (II) wherein X represents $S(O)_2$ and L^1 represents chloro are conveniently prepared by oxidative chlorination of alkyl or benzyl thioethers of formula (IV) (Griffith, O.: J. Biol. Chem., 1983, 258, 3, 1591).

20

21




(IV)

wherein R represents a C1 to 6 alkyl or benzyl residue. Typically R represents unsubstituted benzyl (Ph-CH₂) or tert-butyl.

5

Compounds of formula (IV) may be prepared by reacting a compound of formula (V) in which L² is a leaving group, for example, halo or a sulfonate ester,



(V)

10

with an alkyl or benzyl thiol, R-SH. The reactions are preferably performed in the presence of a base such as diethylisopropylamine or caesium carbonate and in the presence of a suitable solvent, for example, DMF.

15

Compounds of formula (V) may be prepared from, for example, corresponding carboxylic acids and derivatives thereof, using, for example, methods that will be readily apparent to

the man skilled in the art. See, for example, B. George *et al*, J. Org. Chem. 1976, 41(20), 3233; H-C Huang *et al*, J. Med. Chem. 1993, 36(15), 2172; C.J. Crowden *et al*, Tetrahedron Letters, 2000, 41, 8661; Y. Xu *et al*, J. Med. Chem. 2003, 46(24), 5121).

5 It will be appreciated by those skilled in the art that in the processes of the present invention certain potentially reactive functional groups such as hydroxyl or amino groups in the starting reagents or intermediate compounds may need to be protected by suitable protecting groups. Thus, the preparation of the compounds of the invention may involve, at various stages, the addition and removal of one or more protecting groups.

10

Suitable protecting groups and details of processes for adding and removing such groups are described in 'Protective Groups in Organic Chemistry', edited by J.W.F. McOmie, Plenum Press (1973) and 'Protective Groups in Organic Synthesis', 3rd edition, T.W. Greene and P.G.M. Wuts, Wiley-Interscience (1999).

15

The compounds of the invention and intermediates thereto may be isolated from their reaction mixtures and, if necessary further purified, by using standard techniques.

20

The present invention will now be further explained by reference to the following illustrative examples.

25

In the Examples, ¹H-NMR and ¹³C-NMR spectra were recorded on either a Varian Unity Inova 400MHz or Varian Mercury-VX 300MHz instrument. The central solvent peak of dimethylsulfoxide-d₆ (δ_H 2.50 ppm) or tetrahydrofuran-d₈ (δ_H 3.58, 1.73 ppm) were used as internal references. Low resolution mass spectra were obtained on a Agilent 1100 LC-MS system equipped with an APCI ionization chamber.

All solvents and commercial reagents were laboratory grade and used as received. Non-commercially available reagents were synthesised using known literature procedures.

Abbreviations used include:

30

DIEA N,N-diisopropylethylamine

- DCM dichloromethane
THF tetrahydrofuran
THF-D8 deuterated tetrahydrofuran;
AcOH acetic acid;
5 MeCN acetonitrile;
DMF N,N-dimethylformamide;
EtOAc ethyl acetate;
DMSO dimethyl sulfoxide;
DMSO-D6 deuterated dimethyl sulfoxide;
10 Et₂O diethylether;
Et₂NH diethylamine;
TFA trifluoroacetic acid;
IPA 2-propanol;
LC/MS liquid chromatography/mass spectrometry;
15 TLC thin layer chromatography;

Example 1

5-[(4-[(5-Chloropyridin-2-yl)oxyl]piperidin-1-yl)sulfonyl)methyl]-2,4-dihydro-3H-1,2,4-triazol-3-one

20

a) 5-[(Benzylthio)methyl]-2,4-dihydro-3H-1,2,4-triazol-3-one

25

30

Benzylmercaptan (1.75 mL; 14.9 mmol) was dissolved in DMF (20 mL) and solid K₂CO₃ (2.35 g; 17 mmol) was added. To the resulting slurry was added a solution of 5-(chloromethyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (2.0 g; 15 mmol) in DMF (12 mL), prepared by a literature procedure (Eva M. Campi *et. al.*, Tetrahedron, 1994, 50, 8, 2533-2542). The reaction mixture was stirred at room temperature for 20.5 h. Water (80 mL) was added and a thick slurry was formed. The solid product was collected by filtration and washed with water. The remaining filtrate and wash liquid still contained product and was extracted with EtOAc (4 x), the organic phase was then washed with water (2 x), brine (2 x) and dried (Na₂SO₄). Evaporation of solvents gave another crop of crude product. The combined solid materials were suspended in toluene and evaporated to remove water

residues. The crude product was then suspended in a boiling mixture of EtOAc/Heptane (1:4) and allowed to cool before the solid product was collected by filtration. The subtitle compound was obtained as a colourless solid (2.03 g; 61% yield).

APCI-MS m/z : 222.1 $[MH^+]$.

5 1H -NMR (DMSO- D_6): δ 11.35 (1H, vbrs), 11.26 (1H, brs), 7.37-7.21 (5H, m), 3.72 (2H, s), 3.36 (2H, s) ppm.

^{13}C -NMR (DMSO- D_6): δ 156.09, 144.75, 137.66, 128.83, 128.23, 126.79, 34.75, 25.80 ppm.

10 b) (5-Oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methanesulfonyl chloride

5-[(Benzylthio)methyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (0.5 g; 2.26 mmol) was dissolved in AcOH (18 mL) and water (2 mL). The solution was cooled on a ice-bath and Cl_2 gas was slowly bubbled through the solution for 5 min. The green-yellow solution was stirred for 10 min while reaching room temperature and argon gas was bubbled through the solution to remove excess Cl_2 . The clear solution was evaporated to leave an oil which was re-suspended in toluene and evaporated. This process was repeated one more time. The crude product of (5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methanesulfonyl chloride was obtained as a sticky oil still containing benzyl acetate and solvent residues as impurities. This material was dissolved in THF and used directly without further purification.

20 A sample for analytical purposes was obtained by triturating the crude material with isohexane, $CHCl_3$ and Et_2O in that order. After drying under reduced pressure the subtitle compound was obtained as a slightly yellow solid.

1H -NMR (THF- D_8): δ 10.93 (1.4H, vbrs, NH), 5.21 (2H, s, CH_2), 4.80-3.65 (0.9H, vbrs, H_2O + NH) ppm.

25 The reactivity of the sulfonyl chloride was confirmed by its reaction with Et_2NH to give the expected N,N-diethyl-1-(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methanesulfonamide. APCI-MS m/z : 235.1 $[MH^+]$.

30 c) 5-[(4-[(5-Chloropyridin-2-yl)oxy]piperidin-1-yl)sulfonyl)methyl]-2,4-dihydro-3H-1,2,4-triazol-3-one

5-Chloro-2-(piperidin-4-yloxy)pyridine (180 mg; 0.85 mmol) and DIEA (145 μ L; 0.85 mmol) were dissolved in THF (3 mL), and a THF solution of crude (5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methanesulfonyl chloride (approximately 0.56 mmol) was added. The reaction was stirred at room temperature for 1.5 h. Solvent was removed by evaporation and the residue was partitioned between EtOAc and 5% aqueous NaHSO₄ and separated. The water phase was extracted one more time with EtOAc and the combined organic phases were washed with brine and evaporated. The crude product was purified on a preparative HPLC system using a KROMASIL KR-100-7-C18, 250 x 50.8 mm column. A gradient of 20-90% MeCN/water plus 0.1% TFA was used with UV 220 nm for detection. The fractions that according to LC/MS contained the product were evaporated until a slurry was formed and the residual water was removed by freeze drying to leave crude product (40 mg). This material was further purified using a semi-prep HPLC system, KROMASIL 100-5-C18, 250 x 20 mm column, UV 220 nm, and a 80 min gradient of 25-27% MeCN/water plus 50 mM NH₄OAc. Freeze drying gave the title compound as a colourless solid (16 mg; 7.6% yield).
APCI-MS m/z : 374.2 [MH⁺].
¹H-NMR (DMSO-D₆): δ 11.61 (1H, brs), 11.57 (1H, vbrs), 8.20 (1H, d), 7.81 (1H, dd), 6.87 (1H, d), 5.08 (1H, m), 4.32 (2H, s), 3.49-3.39 (2H, m), 3.23-3.13 (2H, m), 2.06-1.94 (2H, m), 1.77-1.64 (2H, m) ppm.
¹³C-NMR (DMSO-D₆): δ 160.79, 155.69, 144.73, 139.14, 137.74, 123.22, 112.74, 69.57, 47.47, 42.80, 30.08 ppm.

Example 2

5-[2-({4-[(5-Chloropyridin-2-yl)oxy]piperidin-1-yl}sulfonyl)ethyl]-2,4-dihydro-3H-1,2,4-triazol-3-one.

a) 5-[2-(Benzylthio)ethyl]-2,4-dihydro-3H-1,2,4-triazol-3-one

3-(Benzylthio)propanoic acid (1.0 g; 5.1 mmol) was dissolved in THF (10 mL). DMF (100 μ L) was added followed by dropwise addition of (COCl)₂ (0.45 mL; 5.2 mmol). After 1 h, a sample for LC was quenched with Et₂NH, showing that approximately 40% starting material remained. More (COCl)₂ (0.12 mL; 1.4 mmol) was added and the reaction mixture

was stirred at room temperature for 2.5 h. A sample for LC was quenched as before with Et₂NH and showed that all starting material had been consumed.

The slightly yellow solution of 3-(benzylthio)propanoyl chloride was added to a pre-cooled solution of semicarbazide hydrochloride (0.95 g; 8.5 mmol) and NaOH (0.83 g; 20.8

5 mmol) in THF (10 mL) and water (2 mL). The slightly acidic (pH 5) solution was neutralised with a few drops of aqueous NaOH to pH 7. The reaction was allowed to reach room temperature and left overnight. A sample was withdrawn for LC/MS analysis and APCI-MS m/z: 254.0 [MH⁺] for the intermediate

2-[3-(benzylthio)propanoyl]hydrazinecarboxamide was found as the major product. To the 10 solution was added 2M aqueous NaOH (30 mL) and the mixture was heated to reflux for 23 h. The reaction mixture was allowed to reach room temperature and acidified using conc. HCl, extracted twice with EtOAc and the organic phase was dried (Na₂SO₄), filtered and evaporated to give crude product (1.08 g). This material was purified using flash chromatography using Si-60 gel and a solvent gradient of 0-10% IPA/DCM. The fractions 15 containing the product were evaporated to give the subtitle compound as colourless solid (0.34 g; 28% yield).

TLC (Si-60, DCM+10%IPA): R_f=0.4.

APCI-MS m/z: 236.1 [MH⁺].

¹H-NMR (DMSO-D₆): δ 11.18 (1H, s), 11.13 (1H, s), 7.36-7.19 (5H, m), 3.75 (2H, s), 20 2.70-2.57 (4H, m) ppm.

b) 2-(5-Oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)ethanesulfonyl chloride

5-[2-(Benzylthio)ethyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (0.3 g; 1.27 mmol) was dissolved in AcOH (18 mL) and water (2 mL). The solution was cooled on an ice/water 25 bath and Cl₂ (g) was slowly bubbled through the stirred solution. When the solution turned greenish yellow the introduction chlorine was stopped. The cold bath was removed and the mixture was stirred for 10 min. Argon (g) was passed through the solution until it became colourless. The clear solution was freeze dried to give the subtitle compound (0.26 g; 97% yield) as a colourless solid.

30 ¹H-NMR (THF-D₈): δ 10.69 (1H, vbrs), 10.57 (1H, brs), 4.29 (2H, m), 3.15 (2H, m) ppm.

c) 5-[2-({4-[(5-Chloropyridin-2-yl)oxy]piperidin-1-yl}sulfonyl)ethyl]-2,4-dihydro-3H-1,2,4-triazol-3-one

5-Chloro-2-(piperidin-4-yloxy)pyridine (100 mg; 0.47 mmol) and DIEA (80 μ L; 0.47 mmol) were dissolved in THF (3 mL). A solution of 2-(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)ethanesulfonyl chloride (65 mg; 0.31 mmol) in THF (4 mL) was added dropwise at room temperature. The reaction was stirred for 1 h before the solvents were removed by evaporation. The residual material was purified using a preparative HPLC system, column Kromasil, KR-100-7-C18, 250 x 50.8 mm. A 40 minute gradient of 20-90% MeCN/water plus 0.1% TFA was used, and UV 220 nm for detection. Fractions containing the desired product were collected. Evaporation of the solvents gave a slurry from which the residual water was removed by freeze drying to give the title compound (90 mg; 74% yield) as a colourless solid.

APCI-MS m/z : 388.1 [MH^+].

1H -NMR (DMSO- D_6): δ 11.25 (1H, s), 11.24 (1H, s), 8.20 (1H, d), 7.81 (1H, dd), 6.88 (1H, d), 5.11 (1H, m), 3.47-3.39 (2H, m), 3.39 (2H, t), 3.23-3.14 (2H, m), 2.81 (2H, t), 2.06-1.96 (2H, m), 1.77-1.66 (2H, m) ppm.

^{13}C -NMR (DMSO- D_6): δ 160.80, 155.86, 144.68, 144.20, 139.11, 123.19, 112.75, 69.58, 45.20, 42.49, 30.15, 20.87 ppm.

Example 3

5-[3-({4-[(5-Chloropyridin-2-yl)oxy]piperidin-1-yl}sulfonyl)propyl]-2,4-dihydro-3H-1,2,4-triazol-3-one

a) 5-(3-Bromopropyl)-2,4-dihydro-3H-1,2,4-triazol-3-one

This was prepared in a similar way as described for 5-(chloromethyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (ref. Eva M. Campi *et. al.*, Tetrahedron, 1994, 50, 8, 2533-2542). Trimethyl 4-bromo-orthobutyrate (5 g; 22 mmol) and semicarbazide hydrochloride (1.12 g; 10 mmol) were stirred in MeOH for 20 h at room temperature. Evaporation of the solvents gave an oily residue that was treated with toluene and evaporated to remove MeOH residues, at which time a precipitate started to form in the toluene solution. The slurry was cooled on dry-ice and the solid material was removed by filtration and washed with

toluene. The solid material (1.79 g) was suspended in water and neutralized with 5% aqueous NaHCO_3 . The product was then extracted into EtOAc, dried over Na_2SO_4 , filtered and evaporated to give the subtitle compound (1.7 g; 82% yield) as a colourless solid. APCI-MS m/z : 206.0+208.0 [MH^+].

¹H-NMR (DMSO- D_6): δ 11.19 (1H, s), 11.11 (1H, s), 3.56 (2H, t), 2.51 (2H, t), 2.09 (2H, quintet) ppm.

b) 5-[3-(Benzylthio)propyl]-2,4-dihydro-3H-1,2,4-triazol-3-one

Benzylmercaptan (0.9 mL; 7.7 mmol) was dissolved in DMF (10 mL) and K_2CO_3 (1.15 g; 8.3 mmol) was added. 5-(3-Bromopropyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (1.6 g; 7.8 mmol) dissolved in DMF (6 mL) was added and the slurry was stirred for 21 h at room temperature. Water (40 mL) was added and an opaque solution was formed which was extracted with EtOAc (4 x). The organic phase was washed with water (2 x) and brine, dried over Na_2SO_4 , filtered and the solvent removed by evaporation. The residual colourless solid was re-dissolved in hot EtOAc (50 mL) and while stirring heptane (150 to 200 mL) was added to precipitate the desired product. After the slurry reached room temperature the solid was collected by filtration and washed with heptane, dried under reduced pressure at +50 °C for 13 h to constant weight to give the subtitle compound (0.7 g; 36% yield) as a colourless solid.

APCI-MS m/z : 250.1 [MH^+].

¹H-NMR (DMSO- D_6): δ 11.16 (1H, s), 11.07 (1H, s), 7.34-7.20 (5H, m), 3.72 (2H, s), 2.43 (2H, t), 2.39 (2H, t), 1.81 (2H, quintet) ppm.

¹³C-NMR (DMSO- D_6): δ 156.01, 146.46, 138.40, 128.64, 128.16, 126.58, 34.69, 29.61, 25.60, 25.19 ppm.

c) 3-(5-Oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)propane-1-sulfonyl chloride.

5-[3-(Benzylthio)propyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (0.5 g; 2.0 mmol) was dissolved in AcOH (18 mL) and water (2 mL). The solution was cooled on an ice/water bath and Cl_2 (g) was bubbled through the solution until a yellow green solution was obtained. The reaction mixture was stirred for 10 min and then the cold bath was removed, Argon (g) was bubbled through the solution until a clear colourless solution was obtained.

Freeze drying gave the sub-title compound as an oil (0.63 g) containing benzyl acetate and solvent residues as major impurities. This material was dissolved in THF and used directly without further purification.

¹H-NMR (THF-D8): δ 12.00-9.20 (2H, baseline broad), 4.05 (2H, t), 2.71 (2H, t), 2.36 (2H, quintet) ppm.

The presence of reactive sulfonylchloride was confirmed by reacting a small sample of the obtained oil with 5-chloro-2-(piperidin-4-yloxy)pyridine to give the expected 5-[3-({4-[(5-chloropyridin-2-yl)oxy]piperidin-1-yl}sulfonyl)propyl]-2,4-dihydro-3H-1,2,4-triazol-3-one.

APCI-MS m/z: 402.1 [MH⁺].

d) 5-[3-({4-[(5-Chloropyridin-2-yl)oxy]piperidin-1-yl}sulfonyl)propyl]-2,4-dihydro-3H-1,2,4-triazol-3-one.

5-Chloro-2-(piperidin-4-yloxy)pyridine (0.16 g; 0.75 mmol) and DIEA (130 uL; 0.76 mmol) were dissolved in THF (3 mL). A THF solution (4 mL) containing crude 3-(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)propane-1-sulfonyl chloride (maximum 0.5 mmol) was slowly added. The reaction was stirred overnight at room temperature and then the yellow slurry was evaporated. The residual material was suspended in MeCN/water and made acidic using a few drops of TFA. The insoluble product was filtered off and dried under reduced pressure. The title compound (137 mg; 68% yield) was obtained as a colourless solid shown to be 95% pure by HPLC.

APCI-MS m/z: 402.2 [MH⁺].

¹H-NMR (DMSO-D6): δ 11.22 (1H, s), 11.30 (1H, s), 8.20 (1H, d), 7.81 (1H, dd), 6.87 (1H, d), 5.11 (1H, m), 3.43 (2H, m), 3.21-3.08 (4H, m), 2.53 (2H, t), 2.08-1.92 (4H, m), 1.72 (2H, m) ppm.

¹³C-NMR (DMSO-D6): δ 160.81, 155.97, 146.04, 144.68, 139.10, 123.71, 112.72, 69.71, 47.42, 42.62, 30.17, 24.65, 19.86 ppm.

Following the general method of Example 1 but substituting the appropriate amine intermediate, the compounds of Examples 4 and 5 were prepared:

Example 4

5-([4-(4-Chlorophenyl)piperazin-1-yl]sulfonyl)methyl)-2,4-dihydro-3H-1,2,4-triazol-3-one.

5 APCI-MS m/z: 358.1 [MH⁺].

¹H-NMR (DMSO-D6): δ 11.61 (1H, s), 11.59 (1H, s), 7.26 (2H, d), 6.98 (2H, d), 4.36 (2H, s), 3.34-3.28 (4H, m), 3.22-3.16 (4H, m) ppm.

Example 5

10 5-([4-[(2-Methoxypyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2H)-yl]sulfonyl)methyl)-2,4-dihydro-3H-1,2,4-triazol-3-one

APCI-MS m/z: 377.1 [MH⁺].

15 ¹H-NMR (DMSO-D6): δ 11.56 (2H, s), 8.72 (2H, s), 6.24 (1H, m), 4.37 (2H, s), 3.95 (3H, s), 3.90 (2H, m), 3.35 (2H, t), 2.35 (2H, m) ppm.

¹³C-NMR (DMSO-D6): δ 163.40, 161.15, 155.63, 137.62, 130.98, 117.81, 111.89, 93.41, 82.08, 54.98, 47.47, 44.54, 41.86, 28.86 ppm.

¹⁵N-¹H-correlated NMR showed a cross peak for two different ¹⁵N at 169.9 and 145.7 ppm to the same ¹H signal at 11.56 ppm.

20 Following the general method of Example 2 but substituting the appropriate amine intermediate, the compounds of Examples 6 and 7 were prepared:

Example 6

25 5-(2-[4-(4-Chlorophenyl)piperidin-1-yl]sulfonyl)ethyl)-2,4-dihydro-3H-1,2,4-triazol-3-one.

APCI-MS m/z: 371.2 [MH⁺].

30 ¹H-NMR (DMSO-D6): δ 11.26 (1H, s), 11.24 (1H, s), 7.36 (2H, d), 7.29 (2H, d), 3.70 (2H, m), 3.39 (2H, t), 2.90 (2H, m), 2.82 (2H, t), 2.66 (1H, m), 1.83 (2H, m), 1.59 (2H, m) ppm.

Example 7

5-(2-([4-(4-Chlorophenyl)piperazin-1-yl]sulfonyl)ethyl)-2,4-dihydro-3H-1,2,4-triazol-3-one.

5 APCI-MS m/z: 372.2 [MH⁺].

¹H-NMR (DMSO-D₆): δ 11.24 (1H, s), 11.22 (1H, s), 7.26 (2H, d), 6.98 (2H, d), 3.42 (2H, t), 3.30 (4H, m), 3.20 (4H, m), 2.82 (2H, t) ppm.

Following the general method of Example 3 but substituting the appropriate amine
10 intermediate, the compounds of Examples 8 and 9 were prepared:

Example 8

5-(3-([4-(4-Chlorophenyl)piperidin-1-yl]sulfonyl)propyl)-2,4-dihydro-3H-1,2,4-triazol-3-one.

15

APCI-MS m/z: 385.3 [MH⁺].

¹H-NMR (DMSO-D₆): δ 11.23 (1H, s), 11.14 (1H, s), 7.36 (2H, d), 7.29 (2H, d), 3.70 (2H, m), 3.13 (2H, t), 2.89 (2H, m), 2.67 (1H, m), 2.54 (2H, t), 1.99 (2H, quintet), 1.83 (2H, m), 1.61 (2H, m) ppm.

20

Example 9

5-(3-([4-(4-Chlorophenyl)piperazin-1-yl]sulfonyl)propyl)-2,4-dihydro-3H-1,2,4-triazol-3-one

25 APCI-MS m/z: 386.2 [MH⁺].

¹H-NMR (DMSO-D₆): δ 11.22 (1H, s), 11.12 (1H, s), 7.26 (2H, d), 6.98 (2H, d), 3.29 (4H, m), 3.22 (4H, m), 3.16 (2H, t), 2.53 (2H, t), 1.99 (2H, quintet) ppm.

30

Pharmacological Example**Isolated Enzyme Assays**

- 5 Recombinant human MMP12 catalytic domain may be expressed and purified as described by Parkar A.A. *et al.*, (2000), Protein Expression and Purification, 20:152. The purified enzyme can be used to monitor inhibitors of activity as follows: MMP12 (50 ng/ml final concentration) is incubated for 60 minutes at room temperature with the synthetic substrate Mac-Pro-Cha-Gly-Nva-His-Ala-Dpa-NH₂ in assay buffer (0.1M "Tris-HCl" (trade mark) buffer, pH 7.3 containing 0.1M NaCl, 20mM CaCl₂, 0.020 mM ZnCl and 0.05% (w/v) "Brij 35" (trade mark) detergent) in the presence (5 concentrations) or absence of inhibitors. Activity is determined by measuring the fluorescence at λ_{ex} 320nm and λ_{em} 405nm. Percent inhibition is calculated as follows: % Inhibition is equal to the
- 10 [Fluorescence_{plus inhibitor} - Fluorescence_{background}] divided by the
- 15 [Fluorescence_{minus inhibitor} - Fluorescence_{background}].


A protocol for testing against other matrix metalloproteinases, including MMP9, using expressed and purified pro MMP is described, for instance, by C. Graham Knight *et al.*, (1992) FEBS Lett. 296(3):263-266.

20 The following table shows the IC₅₀ figures (in nanomolar) for a representative selection of the compounds of the examples when tested against MMP12 and MMP9.

Compound of Example No.	Human MMP12 IC ₅₀ (nm)	Human MMP9 IC ₅₀ (nm)
2	65	318
4	18	414

CLAIMS

1. A compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof



(I)

wherein

R^1 and R^2 independently represent H or C1 to 6 alkyl; said alkyl being optionally further substituted by an aryl ring or an aromatic heterocyclic ring containing 1 to 3 heteroatoms independently selected from O, S and N; said aromatic ring being optionally further substituted by halogen, CF_3 , C1 to 4 alkyl or C1 to 4 alkoxy;

Each R^3 and each R^4 independently represents H or C1 to 6 alkyl; said alkyl being optionally further substituted by OH, C1 to 4 alkoxy, C1 to 4 alkylthio, amino, N-alkylamino or N,N-dialkylamino;

or R^3 and R^4 are bonded together so as to form a 3 to 7 membered ring; said ring optionally incorporating one heteroatom selected from O, $S(O)_q$ and N;

m represents an integer 1, 2 or 3;

X represents a group S(O), S(O)₂ or C(=O);

R⁵ represents H or C1 to 6 alkyl; said alkyl being optionally further substituted by halogen,
5 OH or C1 to 6 alkoxy;

Y represents a direct bond;

or Y and R⁵ are bonded together such that the group -NR⁵Y- together represents a 4 to 7
10 membered saturated or partially unsaturated azacyclic ring; said azacyclic ring optionally
incorporating one further heteroatom selected from O, S(O)_n and N; said azacyclic ring
being optionally benzo fused; said azacyclic ring being optionally substituted by C1 to 6
alkyl, C1 to 6 alkoxy or OH;

15 L represents a direct bond;

or L represents O, S(O)_p, C(O), NR⁶, C(O)NR⁶, NR⁶C(O), C2 to 6 alkynyl, C2 to 6
alkenyl, C1 to 6 alkyl, C1 to 6 heteroalkyl or C3 to 6 heteroalkynyl; said alkyl, alkenyl or
alkynyl group being optionally further substituted by halogen, OH or C1 to 6 alkoxy;

20

n, p and q independently represent an integer 0, 1 or 2;

25

G¹ represents a monocyclic, bicyclic, tricyclic or tetracyclic group comprising one, two,
three or four ring structures each of up to 7 ring atoms; each ring structure being
independently selected from cycloalkyl; cycloalkenyl; heterocycloalkyl; unsaturated
heterocycloalkyl; aryl; or an aromatic heterocyclic ring containing 1 to 3 heteroatoms
independently selected from O, S and N; with each ring structure being independently
optionally substituted by one or more substituents independently selected from halogen,
hydroxy, CHO, C1 to 6 alkyl, C1 to 6 alkoxy, halo-C1 to 6 alkoxy, amino, N-alkylamino,

N,N-dialkylamino, alkylsulfonamino, C2 to 6 alkanoylamino, cyano, nitro, thiol, alkylthio, alkylsulfonyl, alkylaminosulfonyl, C2 to 6 alkanoyl, aminocarbonyl, N-alkylamino-carbonyl, N,N-amino-carbonyl;

- 5 wherein any alkyl radical within any substituent may itself be optionally substituted with one or more groups selected from halogen, hydroxy, C1 to 6 alkoxy, halo-C1 to 6 alkoxy, amino, N-alkylamino, N,N-dialkylamino, N-alkylsulfonamino, N-C2 to 6 alkanoylamino, cyano, nitro, thiol, alkylthio, alkylsulfonyl, N-alkylaminosulfonyl, CHO, C2 to 6 alkanoyl, aminocarbonyl,
- 10 N-alkylaminocarbonyl, N,N-dialkylaminocarbonyl and carbamate;

and wherein any alkyl radical is a C1 to 6 alkyl radical;

- and when G^1 is a bicyclic, tricyclic or tetracyclic group, each ring structure is
- 15 independently joined to the next ring structure by a direct bond, by -O-, by C1-6 alkyl, by C1-6 haloalkyl, by C1-6 heteroalkyl, by C2-6 alkenyl, by C2-6 alkynyl, by sulfone, by CO, by NR^7CO , by $CONR^7$, by NR^7 , by S, or by C(OH), or each ring structure is fused to the next ring structure;

- 20 R^6 and R^7 independently represent H or C1 to 6 alkyl;

and when the group $-NR^5Y-$ represents an azacyclic ring and L represents a direct bond, the group G^1 may also be spiro fused to the azacyclic ring;

- 25 2. A compound according to claim 1, wherein X represents $S(O)_2$.
3. A compound according to claim 1, wherein X represents $C(=O)$.

4. A compound according to any one of claims 1 to 3, wherein R^1 and R^2 each represent hydrogen.
5. A compound according to any one of claims 1 to 4, wherein R^3 and R^4 each represent hydrogen.
6. A compound according to any one of claims 1 to 5, wherein R^5 represents hydrogen or C1 to 6 alkyl and Y represents a direct bond.
7. A compound according to any one of claims 1 to 5, wherein the group $-NR^5Y-$ together represents a five or six membered saturated or partially unsaturated azacyclic ring, said azacyclic ring optionally incorporating one further heteroatom selected from O, $S(O)_n$ and N.
8. A compound according to any one of claims 1 to 7 wherein L represents a direct bond, O, C2 to 6 alkynyl, C1 to 6 alkyl, C1 to 6 heteroalkyl or C3 to 6 heteroalkynyl.
9. A compound according to any one of claims 1 to 8, wherein G^1 represents an optionally substituted monocyclic or bicyclic ring structure.
10. A compound according to claim 1 which is selected from the group consisting of:
- 5-[(4-[(5-chloropyridin-2-yl)oxy]piperidin-1-yl)sulfonyl)methyl]-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-[2-({4-[(5-chloropyridin-2-yl)oxy]piperidin-1-yl}sulfonyl)ethyl]-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-[3-({4-[(5-chloropyridin-2-yl)oxy]piperidin-1-yl}sulfonyl)propyl]-2,4-dihydro-3H-1,2,4-triazol-3-one;
- 5-({4-[(4-chlorophenyl)piperazin-1-yl}sulfonyl)methyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;

5-(2-([4-(4-chlorophenyl)piperidin-1-yl]sulfonyl)ethyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;

5-(2-([4-(4-chlorophenyl)piperazin-1-yl]sulfonyl)ethyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;

5-(3-([4-(4-chlorophenyl)piperidin-1-yl]sulfonyl)propyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;


5-(3-([4-(4-chlorophenyl)piperazin-1-yl]sulfonyl)propyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;

5-([4-([2-methoxypyrimidin-5-yl]ethynyl)-3,6-dihydropyridin-1(2H)-yl]sulfonyl)methyl)-2,4-dihydro-3H-1,2,4-triazol-3-one;

and pharmaceutically acceptable salts and solvates thereof.


11. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as defined in claim 1 which comprises:

15 reaction of a compound of formula (II)



(II)

20 wherein R¹, R², R³, R⁴, X and m are as defined in Claim 1 and L¹ represents a leaving group, with a compound of formula (III)



(III)

wherein G^1 , L, Y and R^5 are as defined in Claim 1;

and optionally thereafter forming a pharmaceutically acceptable salt or solvate.

5

12. A pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 10 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

10 13. A process for the preparation of a pharmaceutical composition as claimed in claim 12 which comprises mixing a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as defined in any one of claims 1 to 10 with a pharmaceutically acceptable adjuvant, diluent or carrier.

15 14. A compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 10 for use in therapy.

20 15. Use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 10 in the manufacture of a medicament for use in the treatment of an obstructive airways disease.

16. Use according to claim 15, wherein the obstructive airways disease is asthma or chronic obstructive pulmonary disease.

17. A method of treating a disease or condition mediated by MMP12 and/or MMP9 which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 10.

5

18. A method of treating an obstructive airways disease which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 10.


10

101414-1 SE

ABSTRACT

COMPOUNDS

5 The invention provides compounds of formula (I):



(I)

wherein R^1 , R^2 , R^3 , R^4 , R^5 , X , Y , L , G^1 and m have the meanings defined in the
10 specification; processes for their preparation; pharmaceutical compositions containing
them; a process for preparing the pharmaceutical compositions; and their use in therapy.